

EXIST (Energetic X-ray Imaging Survey Telescope)




(on behalf of EXIST Science Working Group: http://EXIST.gsfc.nasa.gov)



Imaging the Hard X-ray Universe in Space/Time

EXIST conducts the deepest all-sky imaging HX Survey

Obscured AGN and accretion history of universe



BeppoSAX on Seyfert 2's: EXIST reaches >30X deeper and full sky

Survey black holes on all scales: BH content of Galaxy

EXIST is the Next Generation GRB Mission

- •Survey Gamma Ray Bursts to the limits (z~10-20; ∆time; ∆energy)


 Formation of first massive stars (Pop III)
- Soft Gamma Repeaters (to Virgo); transients, bursts, pulsars (Galaxy)

EXIST



Key EXIST science: Obscured AGN

ASCA and **BeppoSAX** find highly absorbed Seyfert 2's and likely dominant contribution of absorbed AGN to cosmic x-ray background:



and Chandra deep surveys find blank field and optically-dull AGN



EXIST will find >1-10 obscured AGN/square degree and obtain first all-sky measure of Seyfert 2 to QSO 2 luminosity function

EXIST identifies brightest (all sky) obscured AGN for Con-X


EXIST



Key Science: GRB spectral evolution

GRBs have peak flux at ~100-300 keV but extend to >>500 keV.

- High time resolution spectra constrain source models & beaming
- Broad band spectral lags can constrain z (photometric redshifts):



EXIST will measure faintest GRBs over broadest spectral-temporal range (2-3 GRBs/day; 5-50" positions) as the Next Generation GRB Observatory **EXIST** will locate most distant GRBs for Con-X spectra




EXIST Galactic Surveys/Triggers for Con-X

- Supernova and Nova Rates in Galaxy: Stellar Death Rates: HXT images/spectra
 - First map of entire Galaxy in 68,78 keV lines of ⁴⁴Ti: hidden SNR
 - All sky monitoring/imaging of 478,511 keV fast (~8h) line transients: nova rate
- Accretion onto Black Holes vs. Neutron Stars: SXT & HXT images/spectra
 - Spectral variability: accretion instabilities and jet formation (e.g. μ-QSOs)
 - Survey for persistent HX sources in GMCs: ~10-100 Msun BHs or VMOs?
 - Long-term spin measures of accreting pulsars: magnetic accretion
- Magnetic Fields of Neutron Stars: SXT pulsations, spectra
 - Cyclotron lines in LMXBs vs. HMXBs: field decay?
 - Soft Gamma Ray Repeaters (SGRs) in LOCAL GROUP




Mission Concept

Large Area, Wide Field, Broad Band Coded Aperture Telescopes



One of 2 EXIST telescopes: 2 x 2 array of 40° telescopes for combined FOV of 80° x 80°. CZT detector array and CsI shields/collimator shown.



Back -to - back 80° x 80° telescopes mounted at P3 site above truss on ISS for combined 160° x 80° FOV

~8m² imaging CZT, full-sky imaging each orbit >30-300X more sensitive survey at ~5-600keV; ~5-50" GRB & source positions



EXIST Overview

Mission Parameters (low -- high energy)

Energy range (resolution): 5-100 (1) keV; 100 - 600 (3) keV

FOV & resolution: 160° x 80°; 2' - 6'

Sensitivity (5σ): 2mCrab --> 20mCrab/orbit


Temporal resolution: 1µsec --> ~30min; 90min -->1 year

Telescopes/detectors: coded aperture/8m² CZT



EXIST Exposure & Sensitivities

Exposure uniformity (6mo.)



Continuum and line sensitivities (6mo.)





Progress over last year

Captured the interest of the broad astrophysics community


- •Recommended as Moderate Mission in Decadal Survey
- Expanded EXIST Science Working Group
- Presentations at meetings (HEAD, Gamma2001, etc.)

Mission concept engineering study (GSFC/ISAL) for ISS

- Telescope and detector design concept
- Initial ISS interface & background studies
- Formulation office established at GSFC

CZT imaging detectors & balloon flight tests

- Small prototype array & ASIC flown ("CZT2")
- Larger area prototypes & new contact studies









Balloon Flight Test of CZT/ASIC Imager Prototype



Harvard (EXITE2/CZT2) - MSFC (HERO) Ft. Sumner, NM, balloon launch Sept. 19, 2000 (26 hr. flight to Colorado mountains!)






CZT2: Prototype CZT Tiled Array


Test of CZT, pixellated array, and ASIC Readout: µEXIST

- eV Products vs. IMARAD CZT crystals (10 x 10 x 5mm)
- 4 x 4 pixel (2.4mm) arrays on each; tiled across flip-chip crystals
- at focus of capillary optic (~1cm²) to attempt direct "image" of Crab
- 40° x 40° passive/plastic shield collimator for EXIST field of view


Preliminary Results:




Am-241 continuous flight calib. spectrum (60 keV)




Gain vs. time (constant)




Background spectra: total (top) and after shield anti-co (bottom). Am-241 calib. spectrum not yet well subtracted below 60 keV.



CZT2 & data computer in press. vessel



eV& IMARAD CZT in collimator (Am-241 calib. source above)




eV/CZT background image






CZT3: Tiled "Large Area (64cm^2)" CZT Imager

- •Fabrication of Imarad CZT 20 x 20 x 5mm crystals
- Development (with GSFC) of Pt contacts on Imarad CZT (low leakage current; Schottky barrier contacts)
- Development (with IDE) of 2 x 2 ASIC motherboard
- Passive/active collimator and shield; flight test May '01?







EXIST (for Con-X) Development Plans

GSFC/IMDC study of Free Flyer vs. ISS options

- Additional study of ISS option and mission trades (Sept. '01)
- •Engineering study of free flyer mission design (Nov. '01?)
- •Cost analysis of free flyer vs. ISS options

 Schedule &cost for launch by 2009 (survey for Con-X)?

Discussions with potential international partners

Continue discussions with Italian, German, UK groups

Fabrication/packaging CZT imaging detectors

- •Continue development of optimum contacts Imarad/Pt & passivation: HXT tech. dev.
- Studies of large-scale assembly/packaging
- •Balloon flight test of prototype science imager (CZT2 or CZT3 with Imarad/Pt)

